

Pion-less Effective Field Theory on Low-Energy Deuteron Electro-Disintegration

Stefan Christlmeier^a

and

Harald W. Griedhammer^{a,b,1}

^a *Institut für Theoretische Physik (T39), Physik-Department,
Technische Universität München, D-85747 Garching, Germany*

^b *Center for Nuclear Studies, Department of Physics,
The George Washington University, Washington DC 20052, USA*

Abstract

In view of its relation to Big-Bang Nucleo-Synthesis and a reported discrepancy between nuclear models and data taken at S-DALINAC, electro-induced deuteron break-up ${}^2\text{H}(e, e'p)n$ is studied at momentum transfer $q < 100$ MeV and close to threshold in the low-energy nuclear Effective Field Theory without dynamical pions, EFT($\not{\pi}$). The result at N²LO for electric dipole currents and at NLO for magnetic ones converges order-by-order better than quantitatively predicted and contains no free parameter. It is at this order determined by simple, well-known observables. Decomposing the triple differential cross-section into the longitudinal-plus-transverse ($L+T$), transverse-transverse (TT) and longitudinal-transverse interference (LT) terms, we find excellent agreement with a potential-model calculation by Arenhövel et al using the Bonn potential. Theory and data also agree well on σ_{L+T} . There is however no space on the theory-side for the discrepancy of up to 30%, $3\text{-}\sigma$ between theory and experiment in σ_{LT} . From universality of EFT($\not{\pi}$), we conclude that no theoretical approach with the correct deuteron asymptotic wave-function can explain the data. Un-determined short-distance contributions that could affect σ_{LT} enter only at high orders, i.e. at the few-percent level. We notice some issues with the kinematics and normalisation of the data reported.

Suggested PACS numbers: 13.40.-f, 21.30.-x, 21.45.Bc, 25.10.+s, 25.30.Fj, 26.35.+c, 27.10.+h

Suggested Keywords: Effective Field Theory, universality, deuteron electro-disintegration, big-bang nucleosynthesis.

¹Corresponding author; Email: hgried@gwu.edu; permanent address: b

1 Introduction

The deuteron is the simplest nucleus, playing the same fundamental rôle in Nuclear Physics as the hydrogen atom in Atomic Physics. Its electromagnetic properties have been studied in great detail both theoretically and experimentally – amongst others in photo-disintegration (see e.g. [1, 2] for reviews) and electro-disintegration (see e.g. [3, 4] and references therein). The latter process has the advantage to allow for an independent variation of energy and momentum transfer. Most experiments have been performed at higher energy transfers and are in good agreement with potential-model calculations. However, one experiment [5, 6] at the S-DALINAC accelerator of TU Darmstadt (Germany) examined in 2002 the triple-differential cross-section for $d(e, e'p)n$ at low momentum transfer (< 60 MeV/c) and close to the breakup threshold, concentrating on the decomposition into the different structure functions. For the longitudinal-transverse interference cross-section σ_{LT} , a discrepancy of about 30% or up to 3 standard-deviations was discovered relative to the prediction by Arenhövel et al. [4, 5, 7]. As the discrepancy is less pronounced with increasing momentum and energy transfer, this does not necessarily contradict other experiments at higher transfers which are in good agreement with the same potential-model calculations, see e.g. [8].

The disagreement raises however a serious question: The same reaction at the real-photon point, i.e. photo-induced low-energy deuteron breakup and recombination $np \leftrightarrow d\gamma$, is the first and very sensitive step in Big-Bang Nucleo-Synthesis BBN. Indeed, both the higher-energy régime of BBN-relevant photo-dissociation and the S-DALINAC experiment on electro-dissociation are most sensitive to the same electric dipole transition amplitude $E1_V$. Due to the great difficulty of very-low-energy experiments, the lack of data for this reaction in the BBN-relevant energy-region $E \in [20; 300]$ keV makes Nuclear Theory at present the sole provider of input for the BBN network codes [9, 10]. While this has also triggered further experiments at S-DALINAC [11] and the High-Intensity γ -ray Facility HI γ S at TUNL [12, 13], the accuracy claimed by theoretical approaches [1, 14–16] is not yet matched. In view of this, the S-DALINAC findings are the more troubling: How reliably does theory understand the simplest nucleus at low energies, if one of the conceptually simplest non-trivial processes seems to disagree with data? Does this reflect deficits in our understanding of the long-range NN force, meson-exchange currents, “off-shell effects” and cutoff-dependence, or even gauge-invariance? The analysis in Ref. [5] notes that the potential-model approach “without meson-exchange, isobar and relativistic currents accounts for 99% of the final result for σ_{LT} , leaving no room for further improvement” and continues: “At present, there exists no explanation of this surprising result in the framework of conventional nuclear theory. It is an open question whether an alternative interpretation can be offered by effective field theory [...]”. In this work, we answer this question by concluding that *no consistent theoretical approach* can accommodate the data of Ref. [5].

We employ pion-less Effective Field Theory EFT($\not{\pi}$) in the variant in which the effective range is re-summed into the two-nucleon propagator [17–23]. EFT($\not{\pi}$) is well-tested in low-energy reactions with up to three nucleons, see e.g. [24–27] for reviews. Even properties of ^4He [28–30] and ^6Li [31] are now studied successfully. In particular, $np \rightarrow d\gamma$ was studied due to its relevance for BBN [15, 32, 33], culminating in a $N^4\text{LO}$ calculation by Rupak [14]

with a theoretical accuracy of $<1\%$. We here study the S-DALINAC data [5] within the same framework, noting that the lowest energy data show the biggest discrepancy and lie well within the range of applicability of EFT(π). At momenta below the pion-mass, probes cannot resolve the long-range part of the NN -potential as nonlocal. Thus, the most general Lagrangean compatible with the symmetries of QCD can be built out of contact interactions between nucleons as the only dynamical effective hadronic low-energy degrees of freedom. EFT(π) is the unique low-energy version of QCD and allows often (as in this case) for simple, closed-form results. The amplitude is expanded in a small, dimension-less parameter $Q = \frac{p_{\text{typ}}}{\Lambda_\pi}$: the ratio between a typical low-momentum scale p_{typ} of the process involved, and the breakdown scale Λ_π , set by the mass m_π of the pion as the lightest particle which is not included as a dynamical effective degree of freedom. This allows one to estimate the theoretical uncertainties which are induced by neglecting higher-order terms in the momentum expansion of all forces. With a typical low-momentum scale set by the inverse deuteron size, one obtains $Q \approx \frac{1}{3}$. Looking at the order-by-order convergence of all applications considered so far, one finds this estimate to be quite conservative and the actual convergence pattern to be much better. Dimensional regularisation is employed in the two-nucleon system to renormalise high-momentum parts of loops, preserving all symmetries at each step. External currents and relativistic effects are included in a systematic, manifestly gauge-invariant way. No ambiguities arise from meson-exchange currents or off-shell effects.

The presentation is organised as follows: After recalling the essentials of EFT(π) in Sect. 2, the cross-section and individual contributions are given for deuteron electro-disintegration up to next-to-next-to-leading order (N²LO) for electric dipole transitions $E1_V$ and up to NLO for magnetic is-vector dipole transitions $M1_V$. Section 3 contains the detailed comparison with data, focusing on: kinematics in the experiment (Sect. 3.1); the full cross-section (Sect. 3.2) and its decomposition into the structure functions (Sect. 3.3); data normalisation (Sect. 3.4); and possible higher-order effects (Sect. 3.5). After discussing an experiment at slightly higher energies [8] in Sect. 3.6, the Conclusions of Sect. 4 are followed by an Appendix with some useful loop integrals and the explicit form of the hadronic matrix elements. Further background and details are available in S. Christlmeier's Diplom thesis [34].

2 Framework and Calculation

2.1 Kinematics and Cross-Section

The kinematic variables of the disintegration process $d(e, e')n$ are illustrated in Fig. 1. Two frames of reference are conventionally used, see e.g. [3, 4, 35].

On the one hand, the leptonic part $e \rightarrow \gamma^* e'$ of the process is conveniently discussed in terms of variables in the laboratory frame (the deuteron rest frame), denoted by the superscript “lab”. The four-momenta are $(E_0^{\text{lab}}, \vec{k}^{\text{lab}})$ for incoming, and $(E_e^{\text{lab}}, \vec{k}^{\text{lab}})$ for outgoing electrons. The scattering angle between the momenta $\vec{k}^{\text{lab}}$ and $\vec{k}'^{\text{lab}}$ of the incoming and




Figure 1: (Colour online) Kinematics of deuteron electro-disintegration, reproduced with kind permission of [5]. The electron kinematics refers to the lab frame, while the proton variables are defined in the centre-of-mass frame of the two-nucleon final state.

outgoing electron is Θ_e^{lab} . The energy and momentum transfer of the virtual photon is

$$E_X^{\text{lab}} := \omega^{\text{lab}} = E_0^{\text{lab}} - E_e^{\text{lab}} \quad , \quad \vec{q}^{\text{lab}} = \vec{k}^{\text{lab}} - \vec{k}'^{\text{lab}} \quad . \quad (2.1)$$

The hadronic part of the process, $\gamma^* d \rightarrow pn$, is on the other hand more conveniently calculated in its own rest frame, i.e. in the *centre-of-mass frame of the outgoing nucleons* (and thus of the virtual photon and deuteron). Variables in this frame carry no superscript. The outgoing proton (neutron) has momentum $\vec{p}$ ($-\vec{p}$) and kinetic energy $\frac{p^2}{2M}$, $p := |\vec{p}|$. The scattering angles in this system are defined by $\vec{p} \cdot \vec{q} = pq \cos \Theta_p$ and $\vec{p} \cdot (\vec{q} \times \vec{k}) \propto \sin \Phi_p$. Thus, Θ_p is the angle between the virtual photon and the proton, and Φ_p is the angle between the scattering plane, spanned by the incoming and outgoing electron momenta, and the reaction plane, spanned by the proton and photon momenta.

A boost along the momentum transfer $\vec{q}$ transforms between the two frames:

$$\beta = \frac{q^{\text{lab}}}{M_d + \omega^{\text{lab}}} \quad , \quad \gamma = \frac{1}{\sqrt{1 - \beta^2}} \quad , \quad (2.2)$$

where $M_d = 2M - B$ is the deuteron mass. Hence, the quantities defined in the lab frame in (2.1) transform to the proton-neutron cm frame as:

$$\omega = \gamma \omega^{\text{lab}} - \beta \gamma q^{\text{lab}} \quad , \quad q = \beta \gamma M_d \quad . \quad (2.3)$$

We choose as the five independent variables of this process the energies E_0^{lab} , E_e^{lab} (or, equivalently, $E_X^{\text{lab}} = E_0^{\text{lab}} - E_e^{\text{lab}}$) and the scattering angle Θ_e^{lab} of the electrons in the lab frame, plus the proton emission angles Θ_p and Φ_p in the proton-neutron cm frame. The momentum of the outgoing proton is

$$p = \sqrt{(\omega - B)M + q^2 \frac{M}{2M_d}} \quad . \quad (2.4)$$

The momentum vectors are therefore parameterised such that $\vec{q}$ defines the z -direction,

$$\vec{q}^{\text{lab}} = (0, 0, 1)q^{\text{lab}} \quad , \quad (2.5)$$

$$\vec{q} = (0, 0, 1)q \quad , \quad (2.6)$$

$$\vec{k}^{\text{lab}} = (\sin \Theta_{0q}^{\text{lab}}, 0, \cos \Theta_{0q}^{\text{lab}})k^{\text{lab}} \quad , \quad (2.7)$$

$$\vec{k}'^{\text{lab}} = (\sin \Theta_{eq}^{\text{lab}}, 0, \cos \Theta_{eq}^{\text{lab}})k'^{\text{lab}} \quad , \quad (2.8)$$

$$\vec{p} = (\sin \Theta_p \cos \Phi_p, \sin \Theta_p \sin \Phi_p, \cos \Theta_p)p \quad , \quad (2.9)$$

with the angles between the photon and the incoming or outgoing electrons translating as

$$\cos \Theta_{0q}^{\text{lab}} = \frac{k^{\text{lab}} - k'^{\text{lab}} \cos \Theta_e^{\text{lab}}}{q^{\text{lab}}} \quad , \quad \cos \Theta_{eq}^{\text{lab}} = \frac{k^{\text{lab}} \cos \Theta_e^{\text{lab}} - k'^{\text{lab}}}{q^{\text{lab}}} \quad . \quad (2.10)$$

Since the transformation between these two frames is given by a boost along $\vec{q}$, it does not affect the azimuthal angle $\Phi_p \equiv \Phi_p^{\text{lab}}$.

The electron momenta are $\vec{k}_{\text{lab}}^2 = E_{0\text{lab}}^2 - m_e^2$ and $\vec{k}'_{\text{lab}}^2 = E_{e\text{lab}}^2 - m_e^2$. While the electron mass $m_e = 0.511$ MeV does not play a rôle for the experiment at hand, we keep in mind applications to corners of phase-space where its effects could be felt, e.g. in back-scattering [11].

The triple-differential cross-section is then derived from the amplitude $\mathcal{A}$:

$$\frac{d^3\sigma}{dE_e^{\text{lab}} d\Omega_e^{\text{lab}} d\Omega_p} = \frac{k'^{\text{lab}} p M_d M^2}{8(2\pi)^5 (M + \frac{p^2}{2M}) \sqrt{(M_d E_0^{\text{lab}} + \vec{q}^{\text{lab}} \cdot \vec{k}^{\text{lab}})^2 - M_d^2 m_e^2}} |\mathcal{A}|^2 \quad (2.11)$$

$$= \frac{d^3}{dE_e^{\text{lab}} d\Omega_e^{\text{lab}} d\Omega_p} (\sigma_L + \sigma_T + \sigma_{LT} \cos \Phi_p + \sigma_{TT} \cos 2\Phi_p) \quad , \quad (2.12)$$

where the spherical angles are $\Omega_e^{\text{lab}} = (\Theta_e^{\text{lab}}, \Phi_e^{\text{lab}} \equiv 0)$ and $\Omega_p = (\Theta_p, \Phi_p)$. In the second line, the dependence on the azimuthal proton emission angle Φ_p is decomposed into the longitudinal-plus-transverse parts $\sigma_L + \sigma_T$ and the interference terms σ_{LT} and σ_{TT} [3, 4, 35]. For any process to contribute to either of the interference terms, it must from (2.9) depend on those components of the outgoing proton momentum $\vec{p}$ which are transverse to the photon momentum $\vec{q}$. This will become important in the analysis later.

The scattering amplitude $\mathcal{A}$ is rewritten in terms of the photon propagator $D_{\mu\nu}^{(\gamma)}$, the hadronic current J_{hadr}^μ to be calculated in EFT($\not{\Lambda}$) below, and the leptonic current l_μ whose contribution to lowest order in the fine-structure constant is easily found:

$$\mathcal{A} = l^\mu D_{\mu\nu}^{(\gamma)} J_{\text{hadr}}^\nu \quad . \quad (2.13)$$

Current conservation implies $q_\mu l^\mu = 0 = q_\mu J_{\text{hadr}}^\mu$ and is at this order in the fine-structure constant α equivalent to gauge invariance.

If the variables of Ref. [5] are interpreted as discussed in Sect. 3.1, the S-DALINAC experiment was performed at two sets of incident electron energies $E_0^{\text{lab}} \in \{50; 85\}$ MeV, a range of photon energies $E_X^{\text{lab}} \in [8; 16]$ MeV, electron scattering angle $\Theta_e^{\text{lab}} = 40^\circ$ and azimuthal angle $\Phi_p = 45^\circ$. This leads to photon momentum transfers $q \in [32; 65]$ MeV and outgoing proton momenta $p \in [74; 106]$ MeV in the proton-neutron cm-frame. These are the relevant external low-momentum scales of the hadronic matrix element.

2.2 Lagrangean and Parameter Fixing

EFT(π) allows one to address the questions raised by the S-DALINAC data in a model-independent, systematic setting with a reliable estimate of the theoretical uncertainties, free of ambiguities, like off-shell effects and cut-off dependence. As the parts of the effective Lagrangean of EFT(π), its power-counting rules and the parameter-fixing necessary for this work have been discussed extensively by Beane and Savage [22], we repeat them here only briefly. The Feynman rules are also given in [34, App. A].

$$\mathcal{L}_N = N^\dagger \left[iD_0 + \frac{\vec{D}^2}{2M} + \frac{e}{2M}(\kappa_0 + \kappa_1\tau_3)\vec{\sigma} \cdot \vec{B} \right] N, \quad (2.14)$$

$$\mathcal{L}_s = -s_a^\dagger \left[iD_0 + \frac{\vec{D}^2}{4M} - \Delta_s \right] s_a - y_s \left[s_a^\dagger N^T P_a^{(1S_0)} N + \text{H.c.} \right], \quad (2.15)$$

$$\begin{aligned} \mathcal{L}_t = & -t_i^\dagger \left[iD_0 + \frac{\vec{D}^2}{4M} - \Delta_t \right] t_i - y_t \left[t_i^\dagger N^T P_i^{(3S_1)} N + \text{H.c.} \right] \\ & - \frac{C_{sd}}{\sqrt{M\rho_d}} \left[\delta_{ix}\delta_{jy} - \frac{1}{3}\delta_{ij}\delta_{xy} \right] \left[t_i^\dagger (N^T \mathcal{O}_{xy,j} N) + \text{H.c.} \right] - \frac{C_Q}{M\rho_d} t_i^\dagger [iD_0, \mathcal{O}_{ij}] t_j, \end{aligned} \quad (2.16)$$

$$\mathcal{L}_{st} = \frac{eL_1}{M\sqrt{r_0\rho_d}} \left[t_i^\dagger s_3 B_i + \text{H.c.} \right]. \quad (2.17)$$

The one-nucleon Lagrangean $\mathcal{L}_N$ of the nucleon iso-doublet $N = \begin{pmatrix} p \\ n \end{pmatrix}$ with isospin-averaged mass $M = 938.9$ MeV consists of two parts: First, the kinetic term with minimal substitution, $D_\mu = \partial_\mu + ieQA_\mu$, where $Q = \frac{1}{2}(1 + \tau_3)$ is the nucleon charge matrix and $\alpha = \frac{e^2}{4\pi} = \frac{1}{137}$ the fine-structure constant. Second, the interaction via the iso-scalar (iso-vector) magnetic moments, $\kappa_0 = 0.44$ ($\kappa_1 = 2.35$) in nuclear magnetons. The spin (iso-spin) Pauli matrices with vector (iso-vector) index $i = 1, 2, 3$ ($a = 1, 2, 3$) are denoted by σ_i (τ_a).

The Lagrangean for the auxiliary di-baryon field s_a (t_i) in the 1S_0 (3S_1) channel is $\mathcal{L}_s$ ($\mathcal{L}_t$) [17–19, 36]. As we probe only the np system, $D_\mu = \partial_\mu + ieA_\mu$ for both di-baryons. The projection operators are $P_a^{(1S_0)} = \frac{1}{\sqrt{8}}\sigma_2\tau_2\tau_a$, $P_i^{(3S_1)} = \frac{1}{\sqrt{8}}\sigma_2\sigma_i\tau_2$. The auxiliary-field parameters are matched to the effective-range expansion of NN scattering such that the effective range is re-summed to all powers. This speeds up convergence and simplifies calculations of some higher-order effects [17–23]:

$$y_s = \frac{\sqrt{8\pi}}{M\sqrt{r_0}}, \quad \Delta_s = \frac{2}{Mr_0} \left(\frac{1}{a_0} - \mu \right), \quad y_t = \frac{\sqrt{8\pi}}{M\sqrt{\rho_d}}, \quad \Delta_t = \frac{2}{M\rho_d} \left(\gamma - \frac{\rho_d}{2}\gamma^2 - \mu \right) \quad (2.18)$$

The parameter μ encodes the linear cut-off divergence of individual diagrams, calculated in the Power Divergence Subtraction scheme (PDS) version of dimensional regularisation [37, 38]. It is imperative for self-consistency that physical amplitudes are independent of μ . This is indeed the case.

For the spin-singlet, the scattering length and effective range of the np system are probed in deuteron electro-disintegration: $a_0 = -23.71$ fm, $r_0 = 2.73$ fm. Parameters for the spin-triplet are determined from the effective-range expansion about the observed real bound

state, namely the deuteron with binding energy $B = 2.225$ MeV (corresponding to a momentum $\gamma = \sqrt{MB} = 45.70$ MeV) and effective range $\rho_d = 1.764$ fm. Therefore, the deuteron wave function shows the correct exponential decay and normalisation Z already at LO in EFT($\not\pi$) [22, 39]:

$$\Psi_{\text{deuteron}}(r \rightarrow \infty) = \sqrt{\frac{Z}{2\pi\rho_d}} \frac{e^{-\gamma r}}{r} \quad \text{with } Z = \frac{\gamma\rho_d}{1 - \gamma\rho_d} \quad (2.19)$$

The last two terms of $\mathcal{L}_t$ parameterise SD -wave mixing in the spin-triplet, with

$$\mathcal{O}_{xy,j} = -\frac{1}{4} \left(\overleftarrow{D}_x \overleftarrow{D}_y P_j^{(3S_1)} + P_j^{(3S_1)} \overrightarrow{D}_x \overrightarrow{D}_y - \overleftarrow{D}_x P_j^{(3S_1)} \overrightarrow{D}_y - \overleftarrow{D}_y P_j^{(3S_1)} \overrightarrow{D}_x \right) \quad , \quad (2.20)$$

$$\mathcal{O}_{ij} = - \left(\vec{D}_i \vec{D}_j - \frac{1}{3} \delta_{ij} \vec{D}^2 \right) \quad (2.21)$$

operators which ensure that $\mathcal{L}_{st}$ is manifestly gauge-invariant. Their strengths are determined from the asymptotic ratio $\eta_{sd} = 0.0254$ of D - and S -wave components of the deuteron wave function, and of the deuteron quadrupole moment $\mu_Q = 0.2859$ fm² [22]:

$$C_{sd} = \frac{6\sqrt{\pi}\eta_{sd}}{\sqrt{M}\gamma^2} \quad , \quad \mu_Q = 2Z \left[y_t \frac{C_{sd}}{\sqrt{M}\rho_d} \frac{M^2}{32\pi} \frac{2}{3\gamma} + \frac{C_Q}{M\rho_d} \right] \quad (2.22)$$

Here, C_{sd} contributes at LO, and C_Q provides a NLO correction of about 50%.

Finally, $\mathcal{L}_{st}$ in (2.17) parameterises transitions between the 1S_0 and 3S_1 channels by a magnetic field acting on a di-baryon. Its strength is determined by the thermal cross-section $\sigma(E = \frac{p^2}{M} = 1.264 \cdot 10^{-8} \text{ MeV}) = (334.2 \pm 0.5) \text{ mb}$ [40] for radiative capture of neutrons on protons $np \rightarrow d\gamma$. At thermal energies, it is dominated by iso-vectorial $M1_V$ transitions. Magnetic-moment interactions are LO, and the parameter L_1 in $\mathcal{L}_{st}$ enters as a NLO correction of about 50%, while electric transitions are irrelevant. The free parameter is thus determined from the thermal cross-section [22]

$$\sigma^{(M1_V)} = \frac{2\alpha Z(p^2 + \gamma^2)^3}{pM^3} \left| \frac{1}{-\frac{1}{a} + \frac{1}{2}r_0p^2 - ip} \right|^2 \left[\kappa_1 \frac{\gamma - \frac{1}{a} + \frac{1}{2}r_0p^2}{p^2 + \gamma^2} + \frac{L_1}{2} \right]^2 \quad (2.23)$$

as $L_1 = -4.41$ fm [34] or $L_1 = -4.0$ fm [22], depending whether terms quadratic in L_1 are kept. Both variants differ by 10% $\lesssim Q$, as expected from L_1 being a higher-order correction.

We now state the power-counting rules of EFT($\not\pi$) in the version in which the effective-range parameters are treated as unnaturally large and thus are kept at LO together with the scattering lengths [22]. The typical low-energy centre-of-mass momentum $\vec{p}$ and kinetic energy $E/2$ of a nucleon is counted as

$$|\vec{p}| \sim \sqrt{ME} \sim \gamma \sim \frac{1}{\rho_d} \sim \frac{1}{r_0} \sim \frac{1}{a_0} \sim \mu \sim Q\Lambda_{\not\pi} \quad . \quad (2.24)$$

Hence, the wave function renormalisation scales as $Z \sim 1$. SD mixing is suppressed by Q^2 with respect to pure S -wave amplitudes because of the two derivatives in (2.16). Finally,

the ratio of the iso-scalar and iso-vector magnetic moments is $\kappa_0/\kappa_1 \approx 1/5 \lesssim Q$. Therefore, κ_0 is neglected as numerically higher order in the following [41].

Since the gauge field is minimally coupled, these rules are simple to extend: The zero-component of the gauge field A_μ scales like an energy, $A^0 \sim Q^2$; its 3-vector components like a momentum, $\vec{A} \sim Q$. As seen in the previous Sub-Section, photon energies for the S-DALINAC kinematics of deuteron electro-disintegration lie in the range $E_X^{\text{lab}} \in [8; 16]$ MeV, corresponding to $E_X \in [6; 14]$ MeV $\sim [Q^2 \Lambda_\pi^2/M; Q^2 \Lambda_\pi]$, photon momenta in the range $q \in [32; 65]$ MeV $\sim Q\Lambda_\pi$, and outgoing proton momenta in the range $p \in [74; 106]$ MeV $\sim Q\Lambda_\pi$, approaching $\Lambda_\pi \approx m_\pi$. Convergence must therefore be checked with care.

2.3 Electric Contributions to N²LO

Electric dipole transition amplitudes $E1_V$ dominate not only the photo-dissociation cross-section at the higher end of the energy range relevant for BBN, but also electro-disintegration for $p \gtrsim 20$ MeV. To N²LO, the hadronic currents have the structure

$$J_{\text{hadr}}^{(E1_V)\mu} = ie\sqrt{Z} \frac{1}{\sqrt{8}} (N_p^\dagger \sigma^i \sigma_2 N_n^*) \epsilon_{(d)}^j J_{ij}^\mu, \quad (2.25)$$

where the proton and neutron spinors $N_{p/n}$ are specified explicitly. The indices on the current J_{ij}^μ indicate the polarisation j of the initial deuteron and the spin i of the pn -triplet final state. Averaging over spins and polarisations, the electric contribution to (2.11) is

$$|\mathcal{A}_{E1_V}|^2 = \frac{(4\pi\alpha)^2}{(\omega^2 - q^2)^2} \frac{Z}{6} [(k_{\text{lab}}'^\mu J_{\mu}^{\dagger ij} k_{\text{lab}}^\nu J_{\nu}^{ij} + (\mu \leftrightarrow \nu)) - (k_{\mu}^{\text{lab}} k_{\text{lab}}'^\mu - m_e^2) J_{\nu}^{\dagger ij} J^{\nu ij}] . \quad (2.26)$$

The three diagrams contributing at leading order, Q^0 , are shown in Fig. 2. In the




Figure 2: (Colour online) The LO electric contributions to the hadronic current. The double line denotes the spin-triplet di-baryon intermediate state. There is no NLO contribution.

last two graphs, the photon couples to the deuteron which is at this order a pure S-wave state. These diagrams are therefore independent of the direction of the outgoing proton momentum $\vec{p}$, and hence via (2.9) of the proton emission angle Φ_p . They do thus not contribute to the longitudinal-transverse and transverse-transverse parts of the cross-section (2.12). In contradistinction, the final-state interaction in diagram (e1) depends on $\vec{p}$ and thus contributes to σ_{LT} and σ_{TT} .




Figure 3: (Colour online) The order- Q^2 (N^2LO) electric contributions. The SD-mixing vertex proportional to C_{SD} is denoted by a circle; further notation as in Fig. 2.

There are no electric contributions at NLO. The first non-zero corrections, listed in Fig. 3, appear at N^2LO . All arise from SD-mixing proportional to C_{SD} in the Lagrangean (2.17). Relative to the LO contributions, these diagrams are suppressed by two more derivatives, i.e. two more powers of $Q \approx \frac{1}{3}$ and thus should contribute at most $(\frac{1}{3})^2 \approx 10\%$ of the LO terms. The error one makes by truncating the series at this order should be $(\frac{1}{3})^3 \approx 4\%$. The convergence is indeed much better because the asymptotic SD-ratio is numerically $\eta_{sd} = 0.0254 \approx Q^3$, i.e. N^3LO . Even at the S-DALINAC-kinematics of photon momenta as large as 78 MeV ($E_0^{\text{lab}} = 50$ MeV, $E_X^{\text{lab}} = 9$ MeV, $\Theta_e^{\text{lab}} = 40^\circ$, $\Phi_p = 45^\circ$), they amount to not more than 1% of the total; see the discussion and figures in Sects. 3.2 and 3.3 as well as Ref. [34] for details. Only the diagrams (sd1-3) and (sd6) contribute to the interference terms σ_{LT} and σ_{TT} via their final-state interaction or via the D -wave component induced into the deuteron and np final-state wave-function.

The resulting electric currents up to N^2LO are detailed in App. A.2. At each order, they are manifestly gauge-invariant and cutoff-independent.

2.4 Magnetic Contributions to NLO

Magnetic dipole transitions $M1_V$ dominate the inverse reaction $np \rightarrow d\gamma$ at the lower end of the energy range relevant for BBN, but are usually small at S-DALINAC kinematics. We therefore include them only to NLO, Fig. 4. The LO part is set by the iso-vector magnetic moment κ_1 ; the only NLO graph, (m4), comes from the singlet-triplet coupling (2.17).

The hadronic currents from magnetic contributions can to this order be written as

$$J_{\text{hadr}}^{(M1_V)k} = e\sqrt{Z}\epsilon^{ijk}\epsilon_{(d)}^i \frac{1}{\sqrt{8}}(N_p^\dagger \sigma_2 N_n^*) J^j \quad , \quad (2.27)$$




Figure 4: (Colour online) The magnetic contributions to the hadronic current at LO (m1-m3) and NLO (m4). The blob denotes a magnetic photon coupling via the iso-vectorial magnetic moment; the square the dibaryon-photon interaction (2.17) proportional to L_1 ; the dashed double-line the spin-singlet di-baryon intermediate state; further notation as in Fig. 2.

and the squared amplitude is after averaging over initial states:

$$|\mathcal{A}_{M1_V}|^2 = \frac{(4\pi\alpha)^2}{(\omega^2 - q^2)^2} \frac{Z}{6} (\delta^{jm}\delta^{kn} - \delta^{km}\delta^{jn}) \quad (2.28)$$

$$\times [k_n'^{\text{lab}} J^{\dagger j} k_k^{\text{lab}} J^m + k_k'^{\text{lab}} J^m k_n^{\text{lab}} J^{\dagger j} + (k_\mu^{\text{lab}} k'^{\text{lab} \mu} - m_e^2) J^{\dagger j} J^m \delta^{kn}]$$

The vertex where a magnetic photon couples to a di-baryon reduces the amplitude largely, although it is formally NLO. Compared to the electric amplitude, the magnetic one is suppressed by two orders of magnitude in the energy régime of the S-DALINAC experiment, except for the vicinity of $\Theta_p = 70^\circ$; see Figs. 5 to 8.

It is not too difficult to see that the only angular dependence in the magnetic amplitude comes from the $\vec{p} \cdot \vec{q}$ -terms in the intermediate nucleon propagators, as the deuteron and outgoing np -system are again pure S-states in these diagrams. Thus, the amplitudes only test the longitudinal part of $\vec{p}$. There is no Φ_p -dependence, and therefore magnetic transitions do at this order not contribute to σ_{LT} and σ_{TT} interference terms (2.12).

The explicit form of the resulting magnetic currents can be found in App. A.3. At each order, they are manifestly gauge-invariant and cutoff-independent.

3 Theory Confronts Data

We now compare the results of EFT(π) to the S-DALINAC data reported in [5,6] and to the potential model calculation by H. Arenhövel et al. which is based on the Bonn-C r -space potential and includes final-state effects, meson exchange currents, isobar configurations and (in our case negligible) relativistic effects [4, 5, 7]. Before we consider the energy- and angular dependence of the triple differential cross-section and its decomposition, we first have to address a subtle kinematical point.

3.1 Kinematics, Again

In Sect. 2.1, we adopted the standard kinematics of electro-disintegration according to which the leptonic part of the cross-section is calculated in the lab frame (deuteron at rest), while

the hadronic part is determined in the proton-neutron cm-frame of the hadronic sub-process $\gamma^* d \rightarrow pn$. In a slight but revealing abuse of language, the latter is often referred to as “the cm frame”. According to a literal reading of Ref. [5], the S-DALINAC experiment would have been analysed in the centre-of-mass frame of the *whole* process, resulting in large deviations between theory and experiment both in shape and normalisation even of the full triple-differential cross-section, see Fig. 5. We take this to be a slip of the tongue and assume that the experiment was analysed using standard kinematics [3, 4, 35].

Furthermore, the variable which labels the energy transferred by the photon in the S-DALINAC experiment is said to be obtained “after transformation to the centre-of-mass system” [5]. However, agreement of the data with both the potential-model and EFT($\not{\pi}$) calculation is found only if the S-DALINAC-variable “ E_X ” refers to the lab frame. The point $E_X^{(\text{pn})-\text{cm}} = 9$ MeV would correspond to $E_X^{\text{lab}} = 10.6$ MeV and a much smaller cross-section, so that the resulting curve would again substantially deviate from data, see Fig. 5. We therefore interpret “ E_X ” in [5] as E_X^{lab} , defined in the rest frame of the deuteron target.

The experimental results are reported in 4 energy bins, $E_X^{\text{lab}} = [8; 10], [10; 12], [12; 14]$ and $[14; 16]$ MeV. We interpret them such that the count-rate was normalised to give the double-differential cross-section per MeV. Thus, we compare the triple-differential cross-section for E_X^{lab} to the double-differential one *per MeV*. The difference between averaging over the 2 MeV-range and taking the mean value of E_X^{lab} can be neglected down to $E_X^{\text{lab}} = 6$ MeV, as shown in the plot; see also [34]. If E_X is interpreted as given in the pn -cm frame *and* the cross-section as given over the whole 2 MeV-bin, the data would consistently over-shoot the theory results by 10 to 20%.

We are grateful to H. Arenhövel [7] for confirming that the potential-model results reported in [5] were determined in the same kinematics used here.

3.2 Total Differential Cross-Section

Comparing in Fig. 6 to the differential cross-sections reported in Ref. [5] at $E_0^{\text{lab}} = 50$ MeV and 85 MeV in several E_X^{lab} -bins confirms our interpretation of the experiment’s kinematical variables. At $E_0^{\text{lab}} = 85$ MeV, additional data at large Θ_p are available and within their (combined statistical and systematical) error-bars compatible with both theoretical approaches. We note excellent agreement with the calculation by Arenhövel et al. and with the data within error-bars. In Fig. 6, a decomposition of the different contributions in EFT($\not{\pi}$) for the smallest and the largest energy and momentum transfer of the S-DALINAC data is also shown. As expected, the LO electric transitions dominate except around the minimum at $\Theta_p \approx 70^\circ$, where NNLO contributions and magnetic transitions play a significant rôle. That means that by taking into account only minimal coupling of photons to nucleon and di-baryon fields, disintegration can be described highly accurately within EFT($\not{\pi}$). SD mixing, being the only correction up to NNLO, contributes even less than estimated by power counting. This good convergence confirms our confidence that our results are reliable. Less-known short-distance contributions play a very minor rôle, see Sect. 3.5.

The convergence plots also reveal an interesting facet: The N²LO contribution has increased slightly for higher energy and momentum transfer, but much less than expected




Figure 5: (Colour online) Examples of the triple-differential cross-section for $E_X^{\text{lab}} = 9 \text{ MeV}$ at $E_0^{\text{lab}} = 85 \text{ MeV}$, compared to S-DALINAC data [5] (with combined statistical and systematical error-bars) and to the result by Arenhövel et al. (squares) [4, 5, 7]. (a): Interpretation of " $E_X = 9 \text{ MeV}$ " as given in the total cm frame, the pn -cm frame, or the lab-frame; (b): EFT(π) at $E_X^{\text{lab}} = 9 \text{ MeV}$ (solid line) and $E_X^{(\text{pn})-\text{cm}} = 9 \text{ MeV}$ (i.e. $E_X^{\text{lab}} = 10.6 \text{ MeV}$, dashed). The dotted curve is the double-differential cross-section per MeV, obtained by integrating over the bin $E_X^{\text{lab}} \in [8; 10] \text{ MeV}$; the difference to the solid line is almost invisible. (c): comparison on a logarithmic scale of the EFT(π) contributions: electric transitions up to LO and N²LO respectively (recall that NLO₁₁ is zero), and magnetic transitions up to NLO.




Figure 6: (Colour online) EFT(π) results for the differential cross-section, compared to data and calculations by Arenhövel et al. (a): data set at smallest energy and momentum transfer, $E_0^{\text{lab}} = 50 \text{ MeV}$, $E_X^{\text{lab}} = 9 \text{ MeV}$. (b): data sets at largest energy and momentum transfer, $E_0^{\text{lab}} = 85 \text{ MeV}$, $E_X^{\text{lab}} = 15 \text{ MeV}$. (e,f): data set at $E_0^{\text{lab}} = 85 \text{ MeV}$, $E_X^{\text{lab}} = \{11; 13\} \text{ MeV}$. Comparisons of EFT(π) contributions are in Figs. (c,d) included for the smallest and largest energy and momentum transfer: electric transitions up to LO and N²LO respectively (NLO is zero), and magnetic transitions up to NLO.

from naïve dimensional analysis. Deviations between the potential-model and EFT(π) results are also slightly more pronounced there, especially at large Θ_p . As typical momenta in the process approach the breakdown-scale Λ_π of EFT(π), the dimension-less parameter $Q = p_{\text{typ}}/\Lambda_\pi$ approaches unity and the expansion is rendered useless. The momentum transfer $q \in [32; 56]$ MeV is comparable to the intrinsic low-momentum scale of the two-nucleon system, namely the deuteron “binding momentum” $\gamma \approx 45$ MeV. The momentum of the final-state proton in the proton-neutron cm frame ranges however from 75 to 106 MeV and hence becomes at the upper end comparable to $\Lambda_\pi \approx m_\pi$. One would have expected this to affect in particular the final-state interaction diagrams and the SD mixing contributions at N²LO, which depend on $\vec{p}$ in the numerator. The observed enhancement is however much smaller. The good convergence of EFT(π) – even for momenta close to the breakdown scale – was found also in many other applications, see e.g. references in [25–27, 42].

3.3 Comparing Structure Functions

The main goal of the S-DALINAC experiment [5] was a decomposition of the contributions of different structure functions in (2.12): the longitudinal-plus-transverse cross-section, and the LT - and TT -interference parts. However, the uncertainties in the data turned out to be too large for a meaningful comparison of σ_{TT} to theory. Studying the interference terms provides information about the impact of final-state interactions, since these can be Φ_p -dependent, as re-iterated in Sect. 2.1. The contributions of the $L+T$, LT and TT terms are shown in Figs. 7 and 8 for $E_0^{\text{lab}} = 85$ MeV and $E_X^{\text{lab}} = \{9; 11; 13; 15\}$ MeV, again compared to data and the potential-model result.

The results are normalised to $d^3(\sigma_L + \sigma_T)/(dE_e^{\text{lab}} d\Omega_e^{\text{lab}} d\Omega_p)$ at $\Theta_p = 0^\circ$ to make the comparison independent of the absolute normalisation of the data. This also mutes the question whether the cross-sections were averaged or summed over each E_X^{lab} -bin. The error of the value to which the data are normalised has been taken into account in all error bars. $\sigma_L + \sigma_T$ dominates, whereas σ_{LT} and σ_{TT} are one and two orders of magnitude smaller, respectively. However, the interference terms become more relevant with increasing energy transfer, reflecting the growing impact of final-state interactions. In each figure, the result for the LO electric transitions in EFT(π) is also given. Recall that in that case, σ_{LT} and σ_{TT} are nonzero only because of the final-state interaction of the proton with the virtual photon after the deuteron breakup, Fig. 2.(e1). The degree to which the $E1_V$ -transition dominates – already discussed in the previous Section – is even more striking for σ_{LT} . Magnetic transitions do not contribute at all to the interference cross-sections, since their amplitudes are independent of Φ_p , see Sect. 2.4. The impact of SD mixing is increasing slightly with E_X^{lab} , but is small for $\sigma_L + \sigma_T$ and σ_{TT} , and almost negligible for σ_{LT} .

The EFT(π) results are again in good agreement with those of Arenhövel et al. Especially σ_{LT} is essentially identical in both calculations. Only in the very small quantity σ_{TT} can a deviation be detected. It is a factor of 4 larger than the difference between the LO and N²LO results, which in turn is one way to estimate higher-order effects. However, we already hinted above that the contributions from SD mixing and magnetic moment interactions are much smaller than estimated by naïve power-counting. Another estimate for possible corrections is




Figure 7: (Colour online) Decomposition of the triple-differential cross-section into (top to bottom) longitudinal-plus-transverse ($L + T$), longitudinal-transverse (LT) and transverse-transverse (TT) parts at $E_0^{\text{lab}} = 85$ MeV, normalised to σ_{L+T} at $\Theta_p = 0$. Left column: $E_X^{\text{lab}} = 9$ MeV; right column: $E_X^{\text{lab}} = 11$ MeV. No data are available for the TT interference cross-section.




Figure 8: (Colour online) As Figure 7, for $E_X^{\text{lab}} = 13$ MeV (left column); $E_X^{\text{lab}} = 15$ MeV (right column).

to set the uncertainty of the EFT(π) calculation at N²LO as $\sim Q^3 = (p_{\text{typ}}/\Lambda_\pi)^3 \approx [3 \dots 10]\%$ of the LO term, assuming a typical momentum $p_{\text{typ}} \sim q \approx [45; 60]$ MeV. This is apt to over-estimate the theoretical uncertainties, as discussed in Sect. 3.5 in conjunction with possible higher-order corrections. In short, all EFT(π) results should be understood with an uncertainty which reflects an estimate of the higher-order effects not considered. A conservative accuracy limit would be $\lesssim 10\%$ in each observable. It is thus safe to conclude that the potential-model and EFT(π) results agree within the theoretical uncertainties.

3.4 How To Resolve Discrepancies with Data?

The accordance of both theoretical approaches with the data, however, is considerably worse. The discrepancy ranges from just over one standard deviation at the largest energy and momentum transfer, to three and more at the smallest energy and momentum transfer. Not only does the calculated longitudinal-transverse interference cross-section deviate by about 30% from the measured one (as pointed out in [5]), but also $\sigma_L + \sigma_T$ at large Θ_p . In particular, it is remarkable that the latter is over-predicted at $E_X^{\text{lab}} = 9$ MeV, but under-predicted at $E_X^{\text{lab}} = 15$ MeV. A look at $\sigma_L + \sigma_T$ at $\Theta_p = 180^\circ$ *before data normalisation* reveals however a reduced discrepancy, see Fig. 9. Only then do data and theory agree on the shape of the curve, except at the smallest value of E_X^{lab} , and an additional normalisation would solve the problem. While the statistical significance of the discrepancy hardly changes, this may indicate un-accounted systematic errors in the data.

Normalising to $\Theta_p = 0$ seems to deteriorate the accord of calculations and data – although it avoids accounting for the absolute normalisation of data. The reason can be seen by comparing Fig. 9.(b) and (c): The cross-sections both at $\Theta_p = 0$ and $\Theta_p = 180^\circ$ display relatively small deviations which go, however, in different directions: The theoretical result e. g. at $E_X^{\text{lab}} = 15$ MeV is a bit too small for $\Theta_p = 180^\circ$, but too large for $\Theta_p = 0$. In dividing these two values, the discrepancy increases.

The significant difference between data on the one hand, and the agreeing theoretical predictions of EFT(π) and Arenhövel et al. on the other hand, continues for the longitudinal-transverse interference cross section σ_{LT} , see Figs. 7.(b,e) and 8.(b,e). The angular dependence of the normalised values is reproduced, but the normalisation differs between one standard deviations at high energy and momentum transfer, and three standard deviations at high energy and momentum transfer.

Figure 10 examines the energy dependence of the discrepancy for σ_{LT} at the minimum around $\Theta_p = 35^\circ$ as function of E_X^{lab} , both before and after normalisation of the data. Again, the situation becomes, at least for higher energies, slightly better when looking at the absolute values. However, a clear deviation remains, and surprisingly becomes larger at smaller E_X^{lab} . The shapes and slopes of the un-normalised energy-dependence of the minima in σ_{LT} do not match. The un-normalised data at high energy are, at least at non-backward angles, compatible with the theoretical approaches within error-bars, but the discrepancy at smaller energies and momentum transfers worsens, see Fig. 11.




Figure 9: (Colour online) E_X^{lab} -dependence of $\sigma_L + \sigma_T$ at $\Theta_p = 180^\circ$, normalised to $\Theta_p = 0^\circ$ (a) and before normalisation (b). Bottom: $\sigma_L + \sigma_T$ at $\Theta_p = 0^\circ$ (to which all data are normalised). All graphs refer to $E_0^{\text{lab}} = 85$ MeV, $\Theta_e^{\text{lab}} = 40^\circ$.

3.5 Discussion of Higher-Order Contributions

In view of these discrepancies, one must investigate to which extend additional interactions could remedy the problem. The EFT(π) result is complete up to N²LO in electric and up to NLO in magnetic transitions, resulting in a conservative accuracy-estimate of $\lesssim 10\%$, as dis-




Figure 10: (Colour online) E_X^{lab} -dependence of σ_{LT} at the minimum around $\Theta_p = 35^\circ$, normalised (left) and absolute values (right), both for $E_0^{\text{lab}} = 85$ MeV, $\Theta_e^{\text{lab}} = 40^\circ$.




Figure 11: (Colour online) E_X^{lab} -dependence of the *absolute* values of σ_{LT} for the extreme cases $E_X^{\text{lab}} = 9$ MeV (left) and 15 MeV (right) at $E_0^{\text{lab}} = 85$ MeV, $\Theta_e^{\text{lab}} = 40^\circ$. Compare to the normalised data in Figs. 7 and 8.

cussed in Sect. 3.3. Including some renormalisation-group invariant sub-set of higher-order contributions provides a third alternative to estimate the uncertainties of the calculation, besides order-by-order convergence and *prima facie* power counting.

However, the results so far provide a strong argument that higher-order corrections cannot solve the problem: Recall first that σ_{LT} is nonzero only because of non-zero transverse components of the momentum $\vec{p}$ of the outgoing proton. If, in spite of the power-counting, some higher order contributions were important for σ_{LT} , they should therefore contribute *more* at higher energies, where $\vec{p}$ is bigger and thus closer to the break-down scale Λ_π . This would lead to larger higher-order corrections with increasing photon energy. This is however at odds with our analysis of Figs. 7 to 10. As fine-tuning could circumvent this general argument, we explicitly consider in the following some N³LO contributions.

Magnetic transitions are negligible at these energies and in these kinematics, as seen in Figs. 5 and 6. Another contribution to electric transitions includes P -wave nucleon-nucleon final state interactions which also enter at N³LO [33]. While they do become stronger with increasing energy and momentum transfer, they are, even at high energies, not even of the same order of magnitude as the N²LO contributions.

Next, one might think that the correction to the deuteron quadrupole moment of about 50% provided by the C_Q -term in (2.16) can solve the problem at N³LO. However, this term is at best comparable in size to the N²LO term from SD mixing. Its contribution will thus be even smaller than the N²LO correction in Figs. 7 and 8. An explicit calculation [34] renders

$$J_{ij}^0 = -2y_t D_t \frac{C_Q}{M\rho_d} \left(q_i q_j - \frac{1}{3} \mathbf{q}^2 \delta_{ij} \right) , \quad \vec{J}_{ij} = 0 . \quad (3.1)$$

The difference to the N²LO result is less than 0.5 % even at the minimum around $\Theta_p = 70^\circ$ at $E_0^{\text{lab}} = 50$ MeV, $E_X^{\text{lab}} = 9$ MeV, where the effect would need to be biggest to remedy the discrepancy. The contribution of this interaction to the interference terms σ_{LT} and σ_{TT} is furthermore zero since (3.1) is independent of Φ_p .

Corrections from relativistic kinematics are negligible. An example for a relativistic effect which is dynamical but has been neglected in our calculation is the spin-orbit interaction first considered in EFT(π) by Chen et al. [43, 44],

$$\mathcal{L}_{so} = iN^\dagger \left[\left(2\kappa_0 - \frac{1}{2} \right) + \left(2\kappa_1 - \frac{1}{2} \right) \tau_3 \right] \frac{e}{8M^2} \vec{\sigma} \cdot (\vec{D} \times \vec{E} - \vec{E} \times \vec{D}) N , \quad (3.2)$$

where $\vec{E}$ is the electric field. This term is suppressed by $p_{\text{typ}}/M \lesssim Q^2$ relative to the magnetic-field term in $\mathcal{L}_N$ (2.14) and thus enters in $M1_V$ transitions at N⁴LO. Chean et al. also demonstrated that it provides substantial angle-dependent contributions to deuteron Compton scattering at energies around $\omega \approx 49$ MeV [44]. Its Feynman diagrams are those of the LO magnetic transitions, Fig. 4.(m1-3), with the magnetic-moment interaction substituted by (3.2). The vector-component of the hadronic current has the same structure as (2.27), and the zero-component can be written as

$$J_{\text{hadr}}^{0(M1_V)} = e\sqrt{Z}\epsilon^{ijk}\epsilon_{(d)}^i \frac{1}{\sqrt{8}} (N_p^\dagger \sigma_2 N_n^*) J_{jk}^0 . \quad (3.3)$$

The analytical results for these currents are reported in [34]. As expected, they give only a small correction to the already negligible NLO result for magnetic transitions. In σ_{L+T} , they lead to a correction never exceeding 10^{-3} of the LO result. Although they are Φ_p -dependent, the relative difference to LO in σ_{LT} is less than 10^{-2} , with 2×10^{-3} at the extrema. In σ_{TT} , the relative difference to LO at the maximum is 10^{-5} and never exceeds 10^{-4} .

We mention in passing also a N⁴LO contribution which can naïvely be incorporated by modifying the di-baryon propagators (A.12/A.25) to include the shape parameters $\rho_1 = 0.389$ fm³ for the 3S_1 channel and $r_1 = -0.48$ fm³ for the 1S_0 channel of NN -scattering. This neglects diagrams which come from gauging the derivative terms in the Lagrangean which correspond to these corrections, so the result is not complete. However, we see that




Figure 12: (Colour online) Comparison to the data of Ref. [8] for σ_{L+T} (a) and σ_{LT} (b), both in the *lab* frame (also for the hadronic variables!) and normalised to σ_{L+T} at $\Theta_p = 0$.

this leads only to minimal modifications of the structure functions. σ_{L+T} is increased by $\lesssim 5\%$ at back-angles, about 20% in the minimum due to fine-tuning, and in general much less. σ_{LT} changes by less than 1% at the extrema, and σ_{TT} only by 0.1%.

Overall, this discussion supports our estimate of the theoretical uncertainty of our results.

3.6 A Higher-Energy Experiment

In Fig. 12, we finally compare the EFT(π) results to an earlier experiment at a slightly higher momentum transfer [8] which reported good accord between data and the potential model calculation by Arenhövel et al.¹ also for the normalised σ_{LT} . Although the involved proton momentum $p = 117$ MeV is even closer to the breakdown scale of EFT(π) than for the S-DALINAC kinematics, we observe very good agreement for both σ_{L+T} and σ_{LT} . It is remarkable that the result describes the data so well in a régime in which EFT(π) may start to become unreliable, but deviates significantly from data at lower momenta.

4 Concluding Questions

In a parameter-free N²LO calculation using EFT(π), a manifestly gauge and renormalisation-group invariant, well-tested formulation of few-nucleon physics with analytical results which are rooted in a systematic, model-independent low-energy expansion of all nuclear forces, with an *a priori* estimate of theoretical uncertainties corroborated by order-by-order convergence, the S-DALINAC data on deuteron electro-disintegration $d(e, e'p)n$ at low energy and momentum transfer [5, 6] cannot be explained.

¹For this experiment, the calculations were performed using the Paris potential; the difference to the Bonn potential calculation in Ref. [4, 5, 7] should not be relevant here.

Since an important aspect of any Effective Field Theory is its universality and resulting model-independence, this leads us to a conclusion beyond the $\text{EFT}(\not{\pi})$ approach: The predictions of models which reproduce or share the input of $\text{EFT}(\not{\pi})$ must agree with its results to within the accuracy of the EFT calculation, in the range where $\text{EFT}(\not{\pi})$ is applicable. The analysis in Sect. 3 showed that the S-DALINAC data at low momentum and energy transfer lie undoubtedly within this régime. Therefore, any self-consistent potential model, irrespective of the detail led treatment of meson-exchange currents, “off-shell effects”, cutoff-dependence, etc. will to the accuracy outlined here agree with the $\text{EFT}(\not{\pi})$ result, if it incorporates the same ingredients: the same deuteron binding energy, the same asymptotic normalisation Z and asymptotic S-to-D wave ratio η_{sd} of the deuteron wave-function, the same scattering length and effective range in the 1S_0 channel, the same iso-vectorial magnetic moment of the nucleon κ_1 and the same total radiative thermal capture cross-section of neutrons on protons. This is nicely confirmed for electro-disintegration at low energies: The potential-model approach by Arenhövel et al. [4, 5, 7] yields essentially the same results as $\text{EFT}(\not{\pi})$, especially for σ_{LT} .

We find furthermore that σ_{L+T} , σ_{LT} and σ_{TT} are all dominated up to the few-percent level by the leading-order electric transition, i.e. by minimally coupling the photon to the nucleon and deuteron. This coupling is sensitive only to the asymptotics of the deuteron wave-function. The statement is strongest for σ_{LT} , which shows the discrepancy: It is dominated up to the 1%-level by only one LO electric process, Fig. 2.(e1), i.e. by the asymptotic normalisation Z of the deuteron wave-function, eq. (2.19). In contradistinction, the discrepancy to the S-DALINAC data amounts to up to 3 standard-deviations or 30%.

These findings suggest a re-analysis of the experiment [45]. We identified some questions concerning the kinematics and systematics of the experimental analysis and caution that the differences could in part arise from a dis-advantageous normalisation of the data. If the discrepancies were confirmed, this would pose a highly non-trivial problem for Nuclear Theory. Only new data for deuteron electro-disintegration near threshold and at low momentum transfers can settle the issue definitively, particularly concerning the decomposition into the contributions of different structure functions. We are however confident to maintain that the validity and error-estimate of the $E1_V$ -part of the photo-dissociation cross-section relevant for Big-Bang Nucleo-Synthesis as calculated in $\text{EFT}(\not{\pi})$ [14, 15, 32, 33] are not in question.

Acknowledgements

We am indebted to D. B. Kaplan for pointing out to us the puzzle of the S-DALINAC data when they were published. P. von Neumann-Cosel provided excellent insight into the experiment. Both he and H. Arenhövel worked closely with us to resolve the reference-frame issue, answering a barrage of questions with great patience. H. Arenhövel provided invaluable clues by his diligence and by the potential-model results he kindly provided at our request. M. Schwamb helped with comments and further observations, and W. Parke with a careful reading of this manuscript. Our thanks also to N. Kaiser and W. Weise for critical

companionship and a careful reading of the manuscript of the Diplom thesis [34] on which this article is based. H.W.G. gratefully acknowledges the Institut für Theoretische Physik (T39) at TU München and S-DALINAC at TU Darmstadt for its kind hospitality. This work was supported in part by the Bundesministerium für Forschung und Technologie, by the Deutsche Forschungsgemeinschaft under contracts GR1887/2-2 and 3-1, by the CAREER-grant PHY-0645498 of the National Science Foundation and by a Department of Energy grant DE-FG02-95ER-40907.

A Currents and Integrals

We now list the analytical results for the hadronic currents described in Sect. 2.3 and 2.4.

A.1 Useful Loop Integrals

The following loop integrals are calculated by the methodology outlined in [46, 47]², using contour integration for the energy part and the PDS subtraction scheme combined with dimensional integration in D spatial dimension [37, 38] to identify divergences, parameterised by μ , and tensorial reduction. The reader may consult [34, App. D] for details.

The fundamental bubble-sum integral is [37, 38]

$$I_0^{(1)}(a) = \left(\frac{\mu}{2}\right)^{3-D} \int \frac{d^D l}{(2\pi)^D} \frac{1}{(\vec{l} + \frac{\vec{q}}{2})^2 + a} \stackrel{\text{PDS}}{=} \frac{1}{4\pi} (\mu - \sqrt{a}) \quad . \quad (\text{A.1})$$

Reference [46] provides

$$I_0^{(2)}(a, b; q) = \int \frac{d^3 l}{(2\pi)^3} \frac{1}{l^2 + a} \frac{1}{(\vec{l} + \frac{\vec{q}}{2})^2 + b} = \frac{1}{2\pi q} \arctan\left(\frac{q}{2(\sqrt{a} + \sqrt{b})}\right) \quad . \quad (\text{A.2})$$

From that, one finds the μ -independent result with one loop momentum in the numerator

$$A_1(a, b; q) = \int \frac{d^D l}{(2\pi)^D} \frac{\vec{l} \cdot \vec{q}}{l^2 + a} \frac{1}{(\vec{l} + \frac{\vec{q}}{2})^2 + b} = I_0^{(1)}(a) - I_0^{(1)}(b) - \left(\frac{q^2}{4} + b - a\right) I_0^{(2)}(a, b; q) \quad (\text{A.3})$$

and the following μ -dependent ones with two powers of loop momenta in the numerator:

$$\begin{aligned} A_2(a, b; q) &= \frac{1}{D-1} \int \frac{d^D l}{(2\pi)^D} \frac{l^2 - \frac{(\vec{l} \cdot \vec{q})^2}{q^2}}{l^2 + a} \frac{1}{(\vec{l} + \frac{\vec{q}}{2})^2 + b} \\ &= \frac{1}{1-D} \frac{1}{q^2} \left[\frac{2-D}{2} I_0^{(1)}(b) - a I_0^{(2)}(a, b; q) + D \frac{\frac{q^2}{4} + b - a}{q^2} A_1(a, b; q) \right] \quad (\text{A.4}) \end{aligned}$$

$$\begin{aligned} B_2(a, b; q) &= \frac{1}{1-D} \frac{1}{q^2} \int \frac{d^D l}{(2\pi)^D} \frac{l^2 - D \frac{(\vec{l} \cdot \vec{q})^2}{l^2}}{l^2 + a} \frac{1}{(\vec{l} + \frac{\vec{q}}{2})^2 + b} \\ &= \frac{1}{1-D} \frac{1}{q^2} \left[\frac{2-D}{2} I_0^{(1)}(b) - a I_0^{(2)}(a, b; q) + D \frac{\frac{q^2}{4} + b - a}{q^2} A_1(a, b; q) \right] \quad (\text{A.5}) \end{aligned}$$

²We use a slightly different notation.

Three powers of loop momenta in the numerator are covered by the integrals

$$\begin{aligned} A_3(a, b; q) &= \frac{1}{(D-1)q^2} \int \frac{d^D l}{(2\pi)^D} \frac{1}{l^2 + a} \frac{1}{(\vec{l} + \frac{\vec{q}}{2})^2 + b} \left[(\vec{q} \cdot \vec{l}) l^2 - \frac{1}{q^2} (\vec{q} \cdot \vec{l})^3 \right] \\ &= \frac{1}{(D-1)q^2} \left[-\frac{q^2}{2} I_0^{(1)}(b) - a A_1(a, b; q) - \frac{1}{q^2} \hat{A}_3(a, b; q) \right] , \end{aligned} \quad (\text{A.6})$$

$$\begin{aligned} B_3(a, b; q) &= \frac{D+2}{D-1} \frac{1}{q^6} \int \frac{d^D l}{(2\pi)^D} \frac{1}{l^2 + a} \frac{1}{(\vec{l} + \frac{\vec{q}}{2})^2 + b} \left[(\vec{q} \cdot \vec{l})^3 - \frac{3q^2}{D+2} (\vec{q} \cdot \vec{l}) l^2 \right] \\ &= \frac{D+2}{D-1} \frac{1}{q^6} \left[\hat{A}_3(a, b; q) - \frac{3q^2}{D+2} \left(-\frac{q^2}{2} I_0^{(1)}(b) - a A_1(a, b; q) \right) \right] , \end{aligned} \quad (\text{A.7})$$

with the aid of

$$\begin{aligned} \hat{A}_3(a, b; q) &= \int \frac{d^D l}{(2\pi)^D} \frac{(\vec{q} \cdot \vec{l})^3}{l^2 + a} \frac{1}{(\vec{l} + \frac{\vec{q}}{2})^2 + b} \\ &= -\frac{aq^2}{D} I_0^{(1)}(a) - \left(\frac{q^4}{4} - \frac{bq^2}{D} \right) I_0^{(1)}(b) \\ &\quad + \left(\frac{q^2}{4} + b - a \right) \left[-\frac{q^2}{2} I_0^{(1)}(b) + \left(\frac{q^2}{4} + b - a \right) A_1(a, b; q) \right] . \end{aligned} \quad (\text{A.8})$$

A.2 Electric Currents up to N²LO

The electric contributions to the hadronic current at LO are labelled as in Fig. 2. The 4-vector contribution is for the first diagram with $\vec{r} := 2\vec{p} - \vec{q}$

$$J_{ij}^{0(\text{e1})} = -2y_t D_p(\vec{p}; \omega, \vec{q}) \delta_{ij} \quad ; \quad \vec{J}_{ij}^{(\text{e1})} = J_{ij}^{0(\text{e1})} \frac{\vec{r}}{2M} , \quad (\text{A.9})$$

where the proton propagator

$$iD_p(\vec{p}; \omega, \vec{q}) = \frac{i}{-\omega + \frac{\vec{p} \cdot \vec{q}}{M} - \frac{q^2}{2M}} ; \quad (\text{A.10})$$

could be approximated for real photons by dropping the q^2 -term in the denominator ($\omega = q$), but not in electro-disintegration because the kinematics imposes $\omega \sim q^2/M$.

The second diagram gives

$$J_{ij}^{0(\text{e2})} = 2y_t D_t(p) \delta_{ij} \quad ; \quad \vec{J}_{ij}^{(\text{e2})} = -J_{ij}^{0(\text{e2})} \frac{\vec{q}}{4M} , \quad (\text{A.11})$$

with the spin-triplet di-baryon propagator

$$iD_t(p) = \frac{M\rho_d}{2} \frac{i}{\gamma - \frac{\rho_d}{2}(\gamma^2 + p^2) + ip} . \quad (\text{A.12})$$

After integrating over the loop, diagram (3) results in

$$J_{ij}^{0(e3)} = -2y_t^3 D_t(p) M^2 I_0^{(2)}(-p^2, \gamma^2) \delta_{ij} , \quad (\text{A.13})$$

$$\vec{J}_{ij}^{(e3)} = y_t^3 D_t(p) M \left[I_0^{(2)}(-p^2, \gamma^2) + \frac{2}{q^2} A_1(-p^2, \gamma^2; q) \right] \vec{q} \delta_{ij} . \quad (\text{A.14})$$

Here, as in the following, $-p^2$ is understood to be $-p^2 - i\epsilon$ with $\epsilon \searrow 0$, i. e. $\sqrt{-p^2} = -ip$, where p is the proton momentum.

The N²LO contributions come from SD-mixing operators only. Following the numeration of Fig. 3, one finds:

$$J_{ij}^{0(\text{sd1})} = -\frac{1}{2} \frac{C_{sd}}{\sqrt{M\rho_d}} D_p(\vec{p}; \omega, \vec{q}) \left(r_i r_j - \frac{1}{3} \vec{r}^2 \delta_{ij} \right) , \quad \vec{J}_{ij}^{(\text{sd1})} = J_{ij}^{0(\text{sd1})} \frac{\vec{r}}{2M} , \quad (\text{A.15})$$

$$J_{ij}^{0(\text{sd2})} = \frac{C_{sd}}{\sqrt{M\rho_d}} D_t(p) \left(p_i p_j - \frac{1}{3} \vec{p}^2 \delta_{ij} \right) , \quad \vec{J}_{ij}^{(\text{sd2})} = -J_{ij}^{0(\text{sd2})} \frac{\vec{q}}{4M} , \quad (\text{A.16})$$

$$J_{ij}^{0(\text{sd3})} = -2 \frac{C_{sd} y_t^2}{\sqrt{M\rho_d}} D_t(p) M^2 I_0^{(2)}(-p^2, \gamma^2; q) \left(p_i p_j - \frac{1}{3} \vec{p}^2 \delta_{ij} \right) \quad (\text{A.17})$$

$$\vec{J}_{ij}^{(\text{sd3})} = \frac{C_{sd} y_t^2}{\sqrt{M\rho_d}} D_t(p) M \left[I_0^{(2)}(-p^2, \gamma^2; q) + \frac{2}{q^2} A_1(-p^2, \gamma^2; q) \right] \left(p_i p_j - \frac{1}{3} \vec{p}^2 \delta_{ij} \right) \vec{q} \quad (\text{A.18})$$

$$J_{ij}^{0(\text{sd4+sd5})} = -2 \frac{C_{sd} y_t^2}{\sqrt{M\rho_d}} D_t(p) M^2 [B_2(-p^2, \gamma^2; q) + B_2(\gamma^2, -p^2; q)] \left(q_i q_j - \frac{1}{3} \vec{q}^2 \delta_{ij} \right) , \quad (\text{A.19})$$

$$\begin{aligned} \left(\vec{J}_{ij}^{(\text{sd4+sd5})} \right)_k &= 2 \frac{C_{sd}}{\sqrt{M\rho_d}} D_t(p) y_t^2 M \left[\frac{1}{2} B_2(-p^2, \gamma^2; q) \left(q_i q_j - \frac{1}{3} \vec{q}^2 \delta_{ij} \right) q_k \right. \\ &\quad + [A_3(-p^2, \gamma^2; q) - A_3(\gamma^2, -p^2; q)] \left(q_i \delta_{jk} + q_j \delta_{ik} - \frac{2}{3} q_k \delta_{ij} \right) \\ &\quad \left. + [B_3(-p^2, \gamma^2; q) - B_3(\gamma^2, -p^2; q)] \left(q_i q_j q_k - \frac{1}{3} q^2 q_k \delta_{ij} \right) \right] . \quad (\text{A.20}) \end{aligned}$$

The last three diagrams contribute only to the three-vector component of the current:

$$\left(\vec{J}_{ij}^{(\text{sd6})} \right)_k = \frac{C_{sd}}{\sqrt{M\rho_d}} \left[-p_i \delta_{jk} + \left(-p_j + \frac{1}{2} q_j \right) \delta_{ik} + \frac{2}{3} \left(p_k - \frac{1}{4} q_k \right) \delta_{ij} \right] \quad (\text{A.21})$$

$$\left(\vec{J}_{ij}^{(\text{sd7+sd8})} \right)_k = \frac{C_{sd}}{\sqrt{M\rho_d}} D_t(p) y_t^2 \frac{M}{4} [I_0^{(1)}(\gamma^2) - I_0^{(1)}(-p^2)] \left(q_j \delta_{ik} + q_i \delta_{jk} - \frac{2}{3} q_k \delta_{ij} \right) \quad (\text{A.22})$$

With these results, it is simple to show that the LO and N²LO electric currents are separately gauge-invariant, $q_\mu J_{ij}^\mu = 0$. It is also noteworthy that the total current at each order is independent of the regularisation parameter μ .

A.3 Magnetic Currents up to NLO

The LO diagrams, numbered as in Fig. 4, contribute

$$\vec{J}^{(m1)} = -y_t D_p(\vec{p}; \omega, \vec{q}) \frac{\kappa_1}{M} \vec{q} \ , \quad \vec{J}^{(m2)} = -y_t D_n(\vec{p}; \omega, \vec{q}) \frac{\kappa_1}{M} \vec{q} \quad (\text{A.23})$$

$$\vec{J}^{(m3)} = -2y_t D_s(p) y_s^2 \kappa_1 M I_0^{(2)}(-p^2, \gamma^2; q) \vec{q} \ . \quad (\text{A.24})$$

with neutron propagator $D_n(\vec{p}; \omega, \vec{q}) = D_p(-\vec{p}; \omega, \vec{q})$ and spin-singlet di-baryon propagator

$$iD_s(p) = \frac{Mr_0}{2} \frac{i}{\frac{1}{a_0} - \frac{r_0}{2}p^2 + ip} \ . \quad (\text{A.25})$$

The current provided by the NLO contribution is

$$\vec{J}^{(m4)} = -2y_s D_s(p) \frac{L_1}{M\sqrt{r_0\rho_d}} \vec{q} \ . \quad (\text{A.26})$$

Notice that these currents are all transversal only.

References

- [1] H. Arenhövel and M. Sanzone, *Photodisintegration of the Deuteron*, Few-Body Syst. Suppl. 3, Springer Verlag 1991.
- [2] R. A. Gilman and F. Gross, J. Phys. G **28** (2002) R37 [arXiv:nucl-th/0111015].
- [3] W. Fabian and H. Arenhövel, Nucl. Phys. A **314** (1979) 253.
- [4] H. Arenhövel, W. Leidemann and E. L. Tomusiak, Eur. Phys. J. A **23** (2005) 147 [arXiv:nucl-th/0407053].
- [5] P. von Neumann-Cosel et al., Phys. Rev. Lett. **88** (2002) 202304.
- [6] Reported at the website of the experiment, http://qclam.ikp.physik.tu-darmstadt.de/eng/exp_deep.htm.
- [7] H. Arenhövel, private communication.
- [8] T. Tamae et al., Phys. Rev. Lett. **59** (1987) 2919.
- [9] K. M. Nollett and S. Burles, Phys. Rev. D **61**, 123505 (2000) [arXiv:astro-ph/0001440].
- [10] R. H. Cyburt, Phys. Rev. D **70**, 023505 (2004) [arXiv:astro-ph/0401091].
- [11] N. Ryezayeva et al., accepted for publication in Phys. Rev. Lett.
- [12] W. Tornow *et al.*, Phys. Lett. B **574** (2003) 8 [arXiv:nucl-ex/0309009].
- [13] H. Weller, private communication.
- [14] G. Rupak, Nucl. Phys. A **678** (2000) 405 [arXiv:nucl-th/9911018].
- [15] S. Ando, R. H. Cyburt, S. W. Hong and C. H. Hyun, Phys. Rev. C **74** (2006) 025809 [arXiv:nucl-th/0511074].
- [16] L. E. Marcucci, K. M. Nollett, R. Schiavilla and R. B. Wiringa, Nucl. Phys. A **777** (2006) 111 [arXiv:nucl-th/0402078].
- [17] P. F. Bedaque and U. van Kolck, Phys. Lett. B **428**, 221 (1998) [arXiv:nucl-th/9710073].
- [18] P. F. Bedaque, H.-W. Hammer and U. van Kolck, Phys. Rev. C **58**, R641 (1998) [arXiv:nucl-th/9802057].
- [19] P. F. Bedaque and H. W. Griesshammer, Nucl. Phys. A **671**, 357 (2000) [arXiv:nucl-th/9907077].
- [20] F. Gabbiani, P. F. Bedaque and H. W. Griesshammer, Nucl. Phys. A **675**, 601 (2000) [arXiv:nucl-th/9911034].

- [21] G. Rupak and X. W. Kong, Nucl. Phys. A **717**, 73 (2003) [arXiv:nucl-th/0108059].
- [22] S. R. Beane and M. J. Savage, Nucl. Phys. A **694** (2001) 511 [arXiv:nucl-th/0011067].
- [23] S. I. Ando and C. H. Hyun, Phys. Rev. C **72** (2005) 014008 [arXiv:nucl-th/0407103].
- [24] U. van Kolck, Prog. Part. Nucl. Phys. **43**, 337 (1999) [arXiv:nucl-th/9902015].
- [25] S. R. Beane, P. F. Bedaque, W. C. Haxton, D. R. Phillips and M. J. Savage, in (ed.) M. Shifman, At the frontier of particle physics, Vol. 1, p. 133, World Scientific, Singapore (2001) [arXiv:nucl-th/0008064].
- [26] D. R. Phillips, Czech. J. Phys. **52** (2002) B49 [arXiv:nucl-th/0203040].
- [27] P. F. Bedaque and U. van Kolck, Ann. Rev. Nucl. Part. Sci. **52** (2002) 339 [arXiv:nucl-th/0203055].
- [28] L. Platter, H. W. Hammer and U. G. Meissner, Phys. Rev. A **70** (2004) 052101 [arXiv:cond-mat/0404313].
- [29] L. Platter, H. W. Hammer and U. G. Meissner, Phys. Lett. B **607** (2005) 254 [arXiv:nucl-th/0409040].
- [30] H. W. Hammer and L. Platter, Eur. Phys. J. A **32** (2007) 113 [arXiv:nucl-th/0610105].
- [31] I. Stetcu, B. R. Barrett and U. van Kolck, Phys. Lett. B **653** (2007) 358 [arXiv:nucl-th/0609023].
- [32] J. W. Chen, G. Rupak and M. J. Savage, Phys. Lett. B **464** (1999) 1 [arXiv:nucl-th/9905002].
- [33] J. W. Chen and M. J. Savage, Phys. Rev. C **60** (1999) 065205 [arXiv:nucl-th/9907042].
- [34] S. Christlmeier, *Electrodisintegration of the Deuteron in Effective Field Theory*, Diplom thesis supervised by H. W. Griesshammer and available (in English) at http://www.physik.tu-muenchen.de/lehrstuehle/T39/T39_files/T39_people_files/DipThesis/christlmeier.pdf, Technische Universität München, December 2004.
- [35] H. Arenhövel, W. Leidemann and E. L. Tomusiak, Phys. Rev. C **46** (1992) 455.
- [36] In EFT, this was first considered by D. B. Kaplan, Nucl. Phys. B **494** (1997) 471 [arXiv:nucl-th/9610052].
- [37] D. B. Kaplan, M. J. Savage and M. B. Wise, Phys. Lett. B **424** (1998) 390 [arXiv:nucl-th/9801034].
- [38] D. B. Kaplan, M. J. Savage and M. B. Wise, Nucl. Phys. B **534** (1998) 329 [arXiv:nucl-th/9802075].

- [39] D. R. Phillips, G. Rupak and M. J. Savage, Phys. Lett. B **473** (2000) 209 [arXiv:nucl-th/9908054].
- [40] A. E. Cox, S. A. R. Wynchank and C. H. Collie, Nucl. Phys. **74** (1965) 497.
- [41] H. W. Griesshammer and G. Rupak, Phys. Lett. B **529** (2002) 57 [arXiv:nucl-th/0012096].
- [42] H. W. Griesshammer, Nucl. Phys. A **744** (2004) 192 [arXiv:nucl-th/0404073].
- [43] J. W. Chen, X. d. Ji and Y. c. Li, Phys. Lett. B **603** (2004) 5 [arXiv:nucl-th/0407019].
- [44] J. W. Chen, X. d. Ji and Y. c. Li, Phys. Lett. B **620** (2005) 33 [arXiv:nucl-th/0408003].
- [45] A. Shevchenko and P. von Neumann-Cosel, private communication.
- [46] M. J. Savage and R. P. Springer, Nucl. Phys. A **686** (2001) 413 [arXiv:nucl-th/9907069].
- [47] S. Fleming, T. Mehen and I. W. Stewart, Nucl. Phys. A **677** (2000) 313 [arXiv:nucl-th/9911001].